

Sabrina Strickland, MD
Hospital for Special Surgery
523 East 72nd Street
2nd Floor
New York NY 10021-4099

ACL/PCL RECONSTRUCTION with MCL Repair GUIDELINES

	WEIGHT BEARING STATUS	POST OP BRACE	RANGE OF MOTION	THERAPEUTIC GOALS	PATIENT'S CHECKLIST
Week 1-4	50% weight bearing with crutches	Yes Locked in full extension Remove while sleeping	Full 0-90 degrees by week 2 Focus on regaining FULL EXTENSION	Emphasize patient compliance to home exercise program and weight bearing precautions/progression Avoid pain with therapeutic exercise & functional activities Quad muscle reeducation Exercises: Quad sets, heel slides, ankle pumps, gravity assisted knee flexion No open chain knee extension.	<ul style="list-style-type: none"> ○ Bandages should be left on through 1st post op appointment. If bloody or suspect infection call the office immediately. ○ Take Aspirin or Xarelto daily x 30 days ○ Bruising and swelling is normal, it may travel to the ankle. Elevate the leg when laying, perform ankle pumps. ○ Start Physical Therapy 1-2 weeks after surgery - should continue through 6 months or as advised by physician. ○ Patients are responsible for understanding the individual insurance coverage for PT.
Week 5-6	Transition to full weight bearing	Locked in full extension when ambulating	Full	Patellar mobilization Stationary bike No open chain knee extension.	
Week 7-14		Functional brace (if needed)	Full	PT/HEP: LE strength progression within parameters from PT/physician. Continue/normalize phase I exercises as appropriate. No open chain knee extension	

Week 15- 24+		Functional brace if needed ACL brace fitted @ 6mo if indicated		Emphasize patient compliance to maintenance strength program Avoid running until adequate strength development and as advised by your PT and/or physician. Avoid pain with therapeutic exercise & functional activities. Week 36+ No high intensity sport involving cutting such as basketball, soccer, skiing until 9 months and/or as advised by your physician.	
--------------------	--	---	--	---	--

Post Op Brace/CyMedica Evive

- You will be prescribed a long brace for immediately after surgery along with crutches. You will wear the brace for 4-6 weeks. This will help protect your knee until your quad muscle gets stronger. We recommend the CyMedica Evive brace with integrated muscle stimulator. This will help reeducate your quad muscle after surgery and improve your function early. The CyMedica Evive brace requires an upfront deposit which is refundable after processed through your insurance. **Medicare does not provide coverage.**

Game Ready Ice Machine

- We recommend rental of this cold and compression unit. It is an efficient and effective means of controlling swelling and pain post operatively and helps avoid dependency on pain medication. You should be contacted by the hospital vendor for rental information including cost.

Driving Restriction

- Left leg surgery: 1 week for automatic cars, 3-6 weeks for manual transmission cars
- Right leg surgery: 6-8 weeks